



# Analysis of key genes and pathways involved in acute lung injury in a mouse model

Q.H. Han<sup>1\*</sup>, N. Han<sup>1\*</sup>, Y.Z. Liu<sup>1</sup>, Q.H. Jin<sup>2</sup>, Q.Y. Lu<sup>1</sup> and Z.C. Li<sup>1</sup>

<sup>1</sup>Department of Emergency and Trauma Surgery,  
East Hospital Affiliated to Tongji University, Shanghai, China

<sup>2</sup>Internal Medicine, Longhua Community Health Center, Shanghai, China

\*These authors contributed equally to this study.

Corresponding authors: Q.Y. Lu / Z.C. Li

E-mail: lunginjuryluli@hotmail.com

Genet. Mol. Res. 13 (2): 4591-4598 (2014)

Received June 6, 2013

Accepted October 22, 2013

Published June 18, 2014

DOI <http://dx.doi.org/10.4238/2014.June.18.1>

**ABSTRACT.** A mouse model of acute lung injury (ALI) was chosen in this study to explore the key genes and pathways involved in the process of ALI with microarray technology. Gene expression microarray data were downloaded from the Gene Expression Omnibus database. Mice from the experimental group were further divided into 6 subgroups, which received octadecenoate treatments for 1, 1.5, 3, 4, 18, and 24 h. Differentially co-expressed genes were screened to uncover the pathogenesis of ALI. Almost all of the differentially co-expressed genes were identified at two times: 1.5 and 3 h. Functional analysis revealed that several inflammation-related pathways were significantly enriched. Ubiquitin-mediated proteolysis, hematopoietic cell lineage, and leukocyte transendothelial migration were enriched at 1.5 h. The B cell receptor signaling pathway, T cell receptor signaling pathway, natural killer cell-mediated cytotoxicity, Fc epsilon RI signaling pathway, and ubiquitin-mediated proteolysis were significantly enriched at 3 h. It could be inferred that ALI initiated at 1.5 h and lasted through 3 h. However, co-expression patterns were not found from 4 h onward. In conclusion, several key genes and pathways implicated

in the development of ALI were found in this study using the mouse model, among which ubiquitin-mediated proteolysis appears to play an important role in the process.

**Key words:** Acute lung injury; KEGG pathway; Mouse model; Inflammation

## INTRODUCTION

Acute lung injury (ALI) is a secondary lung injury caused by inflammation involving a variety of mediators and effector cells. It is characterized by acute hypoxic respiratory insufficiency and exudative lesions in lung imaging (Bernard et al., 1994). ALI may develop into acute respiratory distress syndrome (ARDS), which carries a high mortality rate (40-50%) (Matthay et al., 2003). The longer the risk factors last, the higher the incidence of ARDS (Iribarren et al., 2000). Studies have indicated a close relationship between the inflammatory response and ALI (Dellinger et al., 2004; Calvano et al., 2005). The inflammatory response is an important part of the body's innate immune system. There are both pro- and anti-inflammatory cytokines in the body; pro-inflammatory cytokines are necessary for initiation of the response, whereas anti-inflammatory cytokines control and "calm down" the response. The timing and balance between the two types of cytokines are important for normal function as excessive inflammatory reactions can cause self tissue damage (Gogos et al., 2000).

High-throughput microarray technology is a suitable choice for studying the mechanism of ALI. Wurfel (2007) proposed a new hypothesis of disease pathogenesis in ARDS with microarray technology. Wang et al. (2008) identified pre-elafin as a biomarker in ARDS through a genome-wide expression analysis. However, some shortcomings of this approach exist. For example, Gogos et al. (2000) used microarray data acquired from animal models suffering from both infection and inflammation, which complicates the expression profiling data and influences the analysis results.

In this study, we adopted mouse models treated with octadecenoate (OA) in order to exclude the potential impact of infection on the expression profile. Although the progression of OA-induced ALI is much faster than that of human ALI, they share similar symptoms. We analyzed microarray data collected at a series of time points to track the change in expression levels over time. Considering the fact that genes and proteins interact to exert biological functions (Han et al., 2004; Huang et al., 2007), we not only screened for differentially co-expressed genes (DCGs), but also performed a pathway analysis to better portray the process of ALI.

## MATERIAL AND METHODS

### Expression microarray data of the mouse model

Mice treated with OA were selected as the animal model for ALI in this study. The gene expression microarray data set GSE18712 (Lesur et al., 2010), consisting of 56 samples, was downloaded from the Gene Expression Omnibus (GEO) database; 51 from the experimental group and 5 from the control group. Raw data were collected from the MusV29K platform and were quantified using the Bzscan2 software. NylonArray was applied for the


diagnosis and normalization of the data. The median was determined as the expression level for probes corresponding to the same symbol.

## Modeling process

Twenty milliliters of OA was intravenously administered to female mice in the experimental group, and an equivalent volume of physiological serum was administered to mice in the control group. Samples were collected at a series of times: 1, 1.5, 3, 4, 18, and 24 h. The number of samples in each subgroup is listed in Table 1, and the comparison model is illustrated in Figure 1.

**Table 1.** Number of samples for experimental groups and the control group.

	Control	1 h	1.5 h	3 h	4 h	18 h	24 h
Samples	5	8	11	13	8	5	6



**Figure 1.** Graphic illustration for the comparison model. OA: octadecenoate,  $\Phi$ : control group.

## Screening of DCGs

The DCGL package (Liu et al., 2010; Yu et al., 2011) of R was used to analyze the microarray data and to screen the DCGs. The DCe method was adopted and the whole screening process consisted of two steps: first, differential co-expression relationships (DCL) were determined among genes, and then genes that were sufficiently associated were identified as DCGs based on the binomial model (Yu et al., 2011). The ‘qth’ method was assigned to determine the relationship, and the cutoff value was set to 0.25. The ‘nbins’ was set to 20 and the P value was set to 0.05 in the LFC model (Mutch et al., 2002).

## Pathway-enrichment analysis

To understand the biological functions of the DCGs, EASE (Hosack et al., 2003) (developed by the DAVID Bioinformatics team) was utilized for KEGG pathway-enrichment analysis. The annotation file was downloaded from DAVID. An EASE score  $<0.1$  was regarded as evidence for significant enrichment.

## RESULTS

### DCGs

The subgroups of the experimental group were respectively compared with the control

group to screen for DCGs. With a false-discovery rate  $\leq 0.05$  set as the cutoff value, DCGs were only identified for 1.5 and 3 h, and the values were relatively low. Extending the cutoff to  $P < 0.05$  and  $P < 0.1$ , most of DCGs were still identified at 1.5 and 3 h (Table 2), and no or few DCGs were found at 4 h or later. DCGs screened out on the basis of  $P < 0.1$  were further analyzed with respect to their functions.

**Table 2.** Number of differentially co-expressed genes identified based upon various cutoff values.

	1 h	1.5 h	3 h	4 h	18 h	24 h
FDR < 0.05	0	36	14	0	0	0
P < 0.05	0	178	81	0	0	1
P < 0.1	0	256	206	0	0	2

FDR = false-discovery rate.

### Pathway-enrichment analysis

KEGG pathway analysis was performed for the 256 DCGs identified at 1.5 h. A total of 6 functional terms were significantly enriched at  $P < 0.1$  (Table 3). The most significantly enriched pathway was ubiquitin-mediated proteolysis.

**Table 3.** Functional analysis results for differentially co-expressed genes identified at 1.5 h.

Term	Genes	P
mmu04120:Ubiquitin mediated proteolysis	Rchy1, Mdm2, Cdc20, Pias2, Soes3, Cul7, Cul3, Skp1a, Rbx1	0.003085
mmu04110:Cell cycle	Stag2, Prkdc, Mdm2, Cdc20, Tgfb2, Skp1a, Sfn, Rbx1	0.00826
mmu04640:Hematopoietic cell lineage	Cd5, Il3ra, Mme, H2-Eb1, Gp9, Csf1	0.017703
mmu04670:Leukocyte transendothelial migration	Itgb2, Cldn6, Sip1a, Vcam1, Ezr, Vcl	0.064141
mmu05200:Pathways in cancer	Traf2, Msh6, Wnt5a, Mdm2, Pias2, Jak1, Tgfb2, Lamb1-1, Lef1, Rbx1, Fos	0.066667
mmu03040:Spliceosome	Cherp, Prpf6, Sfsa2, Phf5a, Snrpf, Sfrs3	0.07377

A total of 7 functional terms were significantly enriched for the 206 DCGs identified at 3 h (Table 4). Several terms were found to be associated with inflammation, such as the B cell receptor signaling pathway, T cell receptor signaling pathway, and natural killer cell-mediated cytotoxicity. Ubiquitin-mediated proteolysis remained significant at this time, which may suggest its importance in the inflammatory cascade.

**Table 4.** Functional analysis results for differentially co-expressed genes identified at 3 h.

Term	Genes	P
mmu04662:B cell receptor signaling pathway	Nfkbib, Cd19, Btk, Rac2, Rasgrp3, Nras	0.003327
mmu04720:Long-term potentiation	Itpr1, Calm1, Rps6ka2, Ppp1cc, Nras	0.01175
mmu04660:T cell receptor signaling pathway	Nfkbib, Il10, Lat, Cd247, Nras, Nck1	0.016704
mmu04650:Natural killer cell-mediated cytotoxicity	Rac2, Itgb2, Ifnar1, Lat, Cd247, Nras	0.019035
mmu04664:Fc epsilon RI signaling pathway	Btk, Rac2, Lat, Nras	0.085901
mmu04120:Ubiquitin-mediated proteolysis	Cdc34, Ube2g1, Cul5, Fzr1, Ube2g2	0.094154
mmu04910:Insulin signaling pathway	Calm1, Ptpn1, Ppp1cc, Foxo1, Nras	0.098052

Based on the above analysis results, it can be inferred that the period from the injection of OA to 1.5 h is the initiation stage of inflammation. As more and more pathways become involved, such as the B cell, T cell, and natural killer cell pathways, the inflammatory

response proceeds and results in more abnormal reactions, which ultimately cause substantial lung injury.

## DISCUSSION

In this study, we explored the genes and biological pathways implicated in ALI using a mouse model administered with OA. Through analyzing the differential co-expression patterns, we found that most of the DCGs were identified at 1.5 and 3 h. A biological interpretation of the DCGs was performed, which revealed several enriched pathways. Ubiquitin-mediated proteolysis, hematopoietic cell lineage, leukocyte transendothelial migration, and spliceosomes were enriched at 1.5 h. The B cell receptor signaling pathway, T cell receptor signaling pathway, natural killer cell-mediated cytotoxicity, and the Fc epsilon RI signaling pathway were significantly enriched at 3 h.

Lesur et al. (2010) conducted a histological examination for the ALI mouse model and reported that lung injury proceeds 1.5 h after OA injection along with interstitial neutrophil accumulation and edema; however, the lesions halted and edema disappeared 3 h after injection. We reached similar conclusions according to the differential co-expression patterns. We speculated that there was a peak between 1.5 and 3 h, which showed the maximum number of DCGs. As the stimulus gradually disappeared over time, the tissue restored to the normal state, and thus few DCGs were identified at 4 h and later. These results suggested that the DCGs could play important roles in the initiation and development of ALI.

Functional analysis for the DCGs revealed that ubiquitin-mediated proteolysis was the most significantly enriched pathway. Previous studies have indicated its regulatory role in the immune system response (Natoli and Chiocca, 2008; El-Hashim et al., 2011). Ubiquitination is involved in the development of the immune system and in regular immune responses through affecting the stability or activity of proteins. For example, ubiquitination is closely associated with the activation of nuclear factor-kappa B, which is an important transcription factor and controls the immune response, inflammatory response, and apoptosis process (Chen, 2005; Bhoj and Chen, 2009; Sun, 2012). E3 ubiquitin protein ligases play a big role in the regulation process, and some of them have also been shown to be involved in the inflammatory response, including CBLB (Bachmaier et al., 2007). Murine double-minute 2 (MDM2) and suppressor of cytokine signaling 3 (SOCS3), which were identified in our study, are also associated with inflammation. MDM2 is believed to link inflammation with epithelial healing during acute kidney injury (Mulay et al., 2012), while SOCS3 is able to attenuate pro-inflammatory signaling (Jo et al., 2005; White et al., 2011). Ligases, such as CUL3, CUL5, and CUL7 were also found to be differentially co-expressed. Studies have revealed their association with cancer (Kim et al., 2007; Lubbers et al., 2011; Thu et al., 2011) and embryonic development (Kasper et al., 2006), but their roles in ALI remain to be elucidated. Furthermore, studies have shown that a variety of ubiquitin ligases are involved in preventing the immune system from attacking self tissues, and that the dysfunction of ubiquitin ligase could lead to the development of autoimmune diseases (Deng et al., 2000; Uchida et al., 2004; Bottomley et al., 2005; Bhoj and Chen, 2009).

The hematopoietic cell lineage and leukocyte transendothelial migration were also significantly enriched at 1.5 h. The hematopoietic cell lineage is mainly involved in the development of blood cells, including granulocytes, monocytes, and dendritic cells. Leukocyte

transendothelial migration is also associated with the inflammatory response. When acute inflammation occurs, local tissues produce interleukin (IL)-1 $\beta$ , IL-8, tumor necrosis factor (TNF)- $\alpha$ , and other inflammatory cytokines to up-regulate the adhesion molecules of vascular endothelial cells. Meanwhile, neutrophils adhere to the surface of injured endothelial cells and then transfer into the interstitial and alveolar spaces, thereby releasing a large number of pro-inflammatory mediators such as inflammatory cytokines, peroxides, leukotriene, protease, and platelet-activating factor, and is thus involved in the neutrophil-mediated inflammatory response and lung injury (Dellinger et al., 2008). Migration of leukocytes from the blood into the tissues is important for immunological surveillance and the inflammatory processes. However, if the leukocytes fail to handle the damaged tissues, they will prolong the process and pose damage to self tissues. Indeed, histological observations have shown neutrophil accumulation in the ALI mouse model (Lesur et al., 2010); therefore, it is not surprising that these two pathways were enriched at corresponding time points. A literature search was performed for some of the DCGs involved in these pathways to further understand their roles in the immune response and in ALI. Previous studies have demonstrated that Cd5 and Csf1 are associated with the immune system (Mizoguchi et al., 2003; Loureiro et al., 2008). The protein product of *Vcam1* mediates leukocyte-endothelial cell adhesion and signal transduction (Lobb et al., 1995). Ezrin and Vinculin are also known to play critical roles in cell adhesion, migration, and organization (Hiscox and Jiang, 1999; Poulet et al., 2001; Carisey and Ballestrem, 2011; Shen et al., 2011).

Another unexpected pathway is the spliceosome pathway. The initial pre-mRNA transcription product must go through the process of splicing to form mature mRNA. As many pro- and anti-inflammatory cytokines are produced, more and more spliceosomes are required. Therefore, alternative splicing might also contribute to the change in expression level. Alternative splicing produces a high degree of protein diversity under conditions of low genetic value, which is valuable for immune systems. The DCGs *Prpf6*, *Sf3a2*, and *Phf5a* encode subunits of the spliceosome pathway.

## CONCLUSIONS

Overall, we identified hundreds of DCGs and confirmed their associations with inflammation as well as ALI through functional analysis. These DCGs are potential drug targets, but require further validation to confirm their usefulness.

## ACKNOWLEDGMENTS

Research supported by the Shanghai Medical Key Subject Construction Project (#ZK2012A28), the National Clinical Key Specialty Construction Project and the Pudong New Area of Outstanding Young Medical Talents (#PWRq2012-13).

## REFERENCES

- Bachmaier K, Toya S, Gao X, Triantafillou T, et al. (2007). E3 ubiquitin ligase Cblb regulates the acute inflammatory response underlying lung injury. *Nat. Med.* 13: 920-926.
- Bernard GR, Artigas A, Brigham KL, Carlet J, et al. (1994). Report of the American-European Consensus conference on acute respiratory distress syndrome: definitions, mechanisms, relevant outcomes, and clinical trial coordination. Consensus Committee. *J. Crit. Care* 9: 72-81.

- Bhoj VG and Chen ZJ (2009). Ubiquitylation in innate and adaptive immunity. *Nature* 458: 430-437.
- Bottomley MJ, Stier G, Pennacchini D, Legube G, et al. (2005). NMR structure of the first PHD finger of autoimmune regulator protein (AIRE1). Insights into autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED) disease. *J. Biol. Chem.* 280: 11505-11512.
- Calvano SE, Xiao W, Richards DR, Felciano RM, et al. (2005). A network-based analysis of systemic inflammation in humans. *Nature* 437: 1032-1037.
- Carisey A and Ballestrem C (2011). Vinculin, an adapter protein in control of cell adhesion signalling. *Eur. J. Cell Biol.* 90: 157-163.
- Chen ZJ (2005). Ubiquitin signalling in the NF-kappaB pathway. *Nat. Cell Biol.* 7: 758-765.
- Dellinger RP, Carlet JM, Masur H, Gerlach H, et al. (2004). Surviving Sepsis Campaign guidelines for management of severe sepsis and septic shock. *Crit. Care Med.* 32: 858-873.
- Dellinger RP, Levy MM, Carlet JM, Bion J, et al. (2008). Surviving Sepsis Campaign: international guidelines for management of severe sepsis and septic shock: 2008. *Intensive Care Med.* 34: 17-60.
- Deng L, Wang C, Spencer E, Yang L, et al. (2000). Activation of the IkappaB kinase complex by TRAF6 requires a dimeric ubiquitin-conjugating enzyme complex and a unique polyubiquitin chain. *Cell* 103: 351-361.
- El-Hashim AZ, Renno WM, Abduo HT, Jaffal SM, et al. (2011). Effect of inhibition of the ubiquitin-proteasome-system and IkappaB kinase on airway inflammation and hyperresponsiveness in a murine model of asthma. *Int. J. Immunopathol. Pharmacol.* 24: 33-42.
- Gogos CA, Drosou E, Bassaris HP and Skoutelis A (2000). Pro- versus anti-inflammatory cytokine profile in patients with severe sepsis: a marker for prognosis and future therapeutic options. *J. Infect. Dis.* 181: 176-180.
- Han JD, Bertin N, Hao T, Goldberg DS, et al. (2004). Evidence for dynamically organized modularity in the yeast protein-protein interaction network. *Nature* 430: 88-93.
- Hiscox S and Jiang WG (1999). Ezrin regulates cell-cell and cell-matrix adhesion, a possible role with E-cadherin/beta-catenin. *J. Cell Sci.* 112 (Pt 18): 3081-3090.
- Hosack DA, Dennis G Jr, Sherman BT, Lane HC, et al. (2003). Identifying biological themes within lists of genes with EASE. *Genome Biol.* 4: R70.
- Huang Y, Li H, Hu H, Yan X, et al. (2007). Systematic discovery of functional modules and context-specific functional annotation of human genome. *Bioinformatics* 23: i222-i229.
- Iribarren C, Jacobs DR Jr, Sidney S, Gross MD, et al. (2000). Cigarette smoking, alcohol consumption, and risk of ARDS: a 15-year cohort study in a managed care setting. *Chest* 117: 163-168.
- Jo D, Liu D, Yao S, Collins RD, et al. (2005). Intracellular protein therapy with SOCS3 inhibits inflammation and apoptosis. *Nat. Med.* 11: 892-898.
- Kasper JS, Arai T and DeCaprio JA (2006). A novel p53-binding domain in CUL7. *Biochem. Biophys. Res. Commun.* 348: 132-138.
- Kim SS, Shago M, Kaustov L, Boutros PC, et al. (2007). CUL7 is a novel antiapoptotic oncogene. *Cancer Res.* 67: 9616-9622.
- Lesur I, Textoris J, Loriod B, Courbon C, et al. (2010). Gene expression profiles characterize inflammation stages in the acute lung injury in mice. *PLoS One* 5: e11485.
- Liu BH, Yu H, Tu K, Li C, et al. (2010). DCGL: an R package for identifying differentially coexpressed genes and links from gene expression microarray data. *Bioinformatics* 26: 2637-2638.
- Lobb RR, Antognetti G, Pepinsky RB, Burkly LC, et al. (1995). A direct binding assay for the vascular cell adhesion molecule-1 (VCAM1) interaction with alpha 4 integrins. *Cell Adhes. Commun.* 3: 385-397.
- Loureiro RM, Monaco KA, Kearney JB, Blickarz-Durand CE, et al. (2008). csf1 is required for early embryonic macrophage development: characterization of the csf1(op)/csf1(op) mutation in ES cell-derived macrophages. *Br. J. Haematol.* 141: 739-742.
- Lubbers J, Lewis S, Harper E, Hledin MP, et al. (2011). Resveratrol enhances anti-proliferative effect of VACM-1/cul5 in T47D cancer cells. *Cell Biol. Toxicol.* 27: 95-105.
- Matthay MA, Zimmerman GA, Esmon C, Bhattacharya J, et al. (2003). Future research directions in acute lung injury: summary of a National Heart, Lung, and Blood Institute working group. *Am. J. Respir. Crit. Care Med.* 167: 1027-1035.
- Mizoguchi A, Mizoguchi E, de Jong YP, Takedatsu H, et al. (2003). Role of the CD5 molecule on TCR gammadelta T cell-mediated immune functions: development of germinal centers and chronic intestinal inflammation. *Int. Immunol.* 15: 97-108.
- Mulay SR, Thomasova D, Ryu M and Anders HJ (2012). MDM2 (murine double minute-2) links inflammation and tubular cell healing during acute kidney injury in mice. *Kidney Int.* 81: 1199-1211.
- Mutch DM, Berger A, Mansourian R, Rytz A, et al. (2002). The limit fold change model: a practical approach for selecting differentially expressed genes from microarray data. *BMC Bioinformatics* 3: 17.

- Natoli G and Chiocca S (2008). Nuclear ubiquitin ligases, NF-kappaB degradation, and the control of inflammation. *Sci. Signal.* 1: e1.
- Pouillet P, Gautreau A, Kadare G, Girault JA, et al. (2001). Ezrin interacts with focal adhesion kinase and induces its activation independently of cell-matrix adhesion. *J. Biol. Chem.* 276: 37686-37691.
- Shen K, Tolbert CE, Guilluy C, Swaminathan VS, et al. (2011). The vinculin C-terminal hairpin mediates F-actin bundle formation, focal adhesion, and cell mechanical properties. *J. Biol. Chem.* 286: 45103-45115.
- Sun SC (2012). Immune receptor signaling: from ubiquitination to NF-kappaB activation. *Cell Mol. Immunol.* 9: 97-98.
- Thu KL, Pikor LA, Chari R, Wilson IM, et al. (2011). Genetic disruption of KEAP1/CUL3 E3 ubiquitin ligase complex components is a key mechanism of NF-kappaB pathway activation in lung cancer. *J. Thorac. Oncol.* 6: 1521-1529.
- Uchida D, Hatakeyama S, Matsushima A, Han H, et al. (2004). AIRE functions as an E3 ubiquitin ligase. *J. Exp. Med.* 199: 167-172.
- Wang Z, Beach D, Su L, Zhai R, et al. (2008). A genome-wide expression analysis in blood identifies pre-elafin as a biomarker in ARDS. *Am. J. Respir. Cell Mol. Biol.* 38: 724-732.
- White GE, Cotterill A, Addley MR, Soilleux EJ, et al. (2011). Suppressor of cytokine signalling protein SOCS3 expression is increased at sites of acute and chronic inflammation. *J. Mol. Histol.* 42: 137-151.
- Wurfel MM (2007). Microarray-based analysis of ventilator-induced lung injury. *Proc. Am. Thorac. Soc.* 4: 77-84.
- Yu H, Liu BH, Ye ZQ, Li C, et al. (2011). Link-based quantitative methods to identify differentially coexpressed genes and gene pairs. *BMC Bioinformatics* 12: 315.